

Galactocentric Distance With the OGLE and Hipparcos Red Clump Stars

Bohdan Paczyński

Princeton University Observatory, Princeton, NJ 08544–1001

e-mail: bp@astro.princeton.edu

Krzysztof Z. Stanek

Harvard-Smithsonian Center for Astrophysics, 60 Garden St., MS20, Cambridge, MA 02174

e-mail: kstanek@cfa.harvard.edu

ABSTRACT

We compare red clump stars with parallaxes known to better than 10% in the Hipparcos catalog and corrected for the interstellar extinction, with the OGLE red clump stars in Baade’s Window also corrected for the interstellar extinction. There are ~ 600 and $\sim 10,000$ such stars in the two data sets, respectively. We find empirically that the average I-band magnitude of red clump stars does not depend on their intrinsic color in the range $0.8 < (V - I)_0 < 1.4$. The red clump luminosity function is well represented by a gaussian with the peak at $M_{I_0,m} = -0.26$, and the dispersion $\sigma_{RC} \approx 0.2$ mag. This allows a single step determination of the distance to the galactic center and gives $R_0 = 8.4 \pm 0.4$ kpc. The number of red clump stars is so large that a formal statistical error is only $\sim 1\%$.

The local stars are relatively blue and have a small color dispersion: $\langle(V - I)\rangle = 1.01$, $\sigma_{(V-I)} = 0.08$, while for the bulge stars $\langle(V - I)_0\rangle = 1.22$, $\sigma_{(V-I)_0} = 0.14$. Presumably, the bulge population has a broader range and a higher average metallicity than the local disk population.

Subject headings: galaxy: center – galaxy: fundamental parameters – galaxy: solar neighborhood – galaxy: stellar content – stars: horizontal-branch

1. Introduction

The center of our Galaxy is at $R_0 \approx 8.0 \pm 0.5$ kpc (Reid 1993). The Hipparcos catalogue of parallaxes (Perryman et al. 1997) will improve this distance measurement. The purpose

of this paper is to present an estimate based on the comparison between the red clump giants measured by Hipparcos and by OGLE (Optical Gravitational Lensing Experiment, cf. Udalski et al. 1992). These stars are the metal rich equivalent of the better known horizontal branch stars, and theoretical models predict that their absolute luminosity only weakly depends on their age and chemical composition (Seidel, Demarque, & Weinberg 1989; Castellani, Chieffi, & Straniero 1992; Jimenez, Flynn, & Kotoneva 1997). Indeed the OGLE color-magnitude diagram of stars in Baade’s Window (Udalski et al. 1993; Paczyński et al. 1994, their Fig.1; Kiraga, Paczyński & Stanek 1997, their Fig.6), and the Hipparcos’ absolute magnitude-color diagram (Perryman et al. 1997, their Fig.3) clearly show how compact the red clump is. In this paper we determine the variance in the absolute I band magnitude to be only ~ 0.2 mag.

Any method of the galactocentric distance determination which is based on stars suffers from at least four problems:

1. The accuracy depends on the absolute magnitude determination for the nearby stars.
2. Interstellar extinction has to be determined for the stars near the galactic center as well as for those near the sun.
3. The masses, ages, and chemical composition may be different for the stars near the galactic center and for their counterparts near the sun.
4. The statistical error is large if the number of objects is small.

The red clump giants are the only type of stars which do not suffer from the fourth problem. In spite of their large number and sound theoretical understanding these stars have seldom been used for the galactic structure studies. However, recently Stanek (1995) and Stanek et al. (1994, 1997) used these stars to map the Galactic bar, and also to map the interstellar extinction in Baade’s Window (Woźniak & Stanek 1996; Stanek 1996). In this paper we compare the absolute magnitudes of ~ 600 nearby red clump stars with accurate (better than 10%) trigonometric parallaxes measured by Hipparcos with the apparent magnitudes of the red clump stars measured by the OGLE in Baade’s Window. This comparison gives the galactocentric distance in a single step. In the last section we discuss various effects which have to be evaluated in order to reduce the systematic error.

2. The method and preliminary result

Inspection of the color-magnitude diagrams based on Hipparcos and OGLE data reveal a strong dependence of the visual magnitude of red clump giants on their color, while their I -band magnitudes reveal no color dependence. Thus, on purely observational grounds, the I -band turns out to be the best in the applications in which the red clump stars are used as standard candles. It is possible that bolometric corrections to the I -band are very small for these moderately cool stars, and theoretical models show only weak dependence of M_{bol} on either age or chemical composition (Seidel, Demarque, & Weinberg 1989; Castellani, Chieffi, & Straniero 1992; Jimenez, Flynn, & Kotoneva 1997).

Using the extinction map of Stanek (1996) with the zero point determined by Gould, Popowski & Terndrup (1997) and by Alcock et al. (1997) we selected the region of Baade’s Window with $A_V < 1.5$, to minimize the effects of a possible error in the ratio A_V/E_{V-I} . The $I_0 - (V - I)_0$ color-magnitude diagram for this part of Baade’s Window is shown in Fig.1. We divided the red clump region into five color bins and we fitted the observed distribution of the stars in each bin with a function

$$n(I_0) = a + b(I_0 - I_{0,m}) + \frac{N_{RC}}{\sigma_{RC}\sqrt{2\pi}} \exp \left[- \left(\frac{I_0 - I_{0,m}}{2\sigma_{RC}} \right)^2 \right], \quad (1)$$

where the first two terms describe a fit to the “background” distribution of the red giant stars, and the gaussian term represents a fit to the red clump itself. The locations of five points with the vertical error bars in Fig.1 correspond to the values of $I_{0,m}$ and their formal 3σ errors for the five color bins. No statistically significant color trend is apparent in $I_{0,m}$.

The Hipparcos based absolute magnitude-color diagram is shown in Fig.2 for the stars with parallaxes measured to better than 10%. The Cousins’ $(V - I)$ colors in the catalog come from various sources. For this study we used only those stars for which the source of $(V - I)$ was labeled A, C, E, F, or G, i.e. those for which I -band photometry was available. There are 664 such stars within the dashed rectangle in Fig.2. We divided them into two color bins and fitted their distribution with the formula analogous to that given in Eq.1, with the I_0 replaced with the M_I . There is no significant color trend in M_I .

Given the fact that the average luminosity of the red clump stars appears to be independent of their color in Baade’s Window (Fig.1) and in the solar neighborhood (Fig.2), it seems safe to determine the distance to the galactic center by comparing these two populations. It is also very fortunate that even though the bulk of the bulge population is redder, and hence it seems to be more metal rich than the solar neighborhood, there is a full overlap in the colors. In this first approximation we neglect the extinction of the Hipparcos red clump giants as their average distance is only 87 pc . We selected the red




Fig. 1.— The color-magnitude diagram corrected for interstellar extinction is shown for the stars in the part of Baade’s Window with a small extinction, $A_V < 1.5$ mag. The dashed rectangle surrounds the red clump region used for the comparison with the local stars. The ridge of the highest density of the red clump stars, $I_{0,m}$, is shown for five color bins in the range $0.8 < (V - I)_0 < 1.4$, as calculated with the Eq.1. It is apparent that the value of $I_{0,m}$ does not depend on the $(V - I)$ color.

clump stars in the color range $0.8 < (V - I)_0 < 1.25$ in Baade’s Window (there were 7174 of them) and $0.8 < (V - I) < 1.25$ in Hipparcos catalog, and we fitted both distributions with Eq.1. We obtained $I_{0,m} = 14.323 \pm 0.009$ and $M_{I,m} = -0.185 \pm 0.016$. Therefore the distance modulus is $(m - M)_{BW} = (I_{0,m} - M_{I,m}) = 14.508 \pm 0.019$, and $d_{BW} = 7.97$ kpc, with the formal statistical error of only ± 80 pc.

The distribution of the two groups of the red clump stars as a function of their absolute I-band magnitude is shown in Fig.3 with two histograms as well as with two analytical fits of the type described by the Eq.1. All these distributions are normalized. The gaussian fitted to the OGLE red clump distribution has $\sigma_{RC} = 0.28$ mag and the gaussian fitted to the Hipparcos distribution has $\sigma_{RC} = 0.24$ mag.

The distribution of the red clump stars as a function of their $(V - I)$ color is shown in Fig.4. It is clear that the bulge population is redder, and presumably more metal rich than the nearby disk population.




Fig. 2.— The color-magnitude diagram is shown for the Hipparcos stars with trigonometric parallax errors smaller than 10%. The dashed rectangle surrounds the red clump region used for the comparison with Baade’s Window stars. The ridge of the highest density of the red clump stars, $M_{I,m}$, is shown for two color bins in the range $0.8 < (V - I) < 1.2$, as calculated with the Eq.1. Note that the average red clump color is $\langle (V - I) \rangle \approx 1.0$, compared to the average value in Baade’s Window $\langle (V - I)_0 \rangle \approx 1.2$, indicating that the galactic bulge stars are substantially more metal rich than the solar neighborhood. Hipparcos $(V - I)$ colors are given to two decimal places only. This figure was made with a random third decimal digit added to the data to avoid the appearance of artificial bands.

3. Correcting some errors

The very small formal statistical error for our galactocentric distance follows from the very large number of red clump stars measured by Hipparcos and OGLE. The number of these stars is several orders of magnitude larger than the number of either RR Lyrae or Mira variables, and the observed dispersion in their magnitudes is only $\sigma_{RC} \approx 0.24 \text{ mag}$. However, in addition to very small statistical errors there are much larger systematic errors, some of which we try to estimate and correct for.

We checked if the 10% error limit adopted by us was not too generous, and we repeated the analysis with a more stringent 5% upper limit to the parallax errors. This reduced the number of stars within the rectangle shown in Fig.2 from 664 to 240, and




Fig. 3.— The number of red clump stars in the solar neighborhood, based on Hipparcos data, is shown as a function of absolute magnitude M_I with the thin solid line (fit and histogram). The number of red clump stars in Baade’s Window, based on the OGLE data, is shown as a function of absolute magnitude M_I with thick solid line, adopting the preliminary distance modulus: $I_0 - M_I = 14.508$. All distributions are normalized.

gave $M_{I,m} = -0.173 \pm 0.027$, well within small statistical errors of $M_{I,m} = -0.185 \pm 0.016$ obtained for the 10% sample. The dispersions were also similar: $\sigma_{RC} = 0.243$ mag for the 10% sample and $\sigma_{RC} = 0.225$ mag for the 5% sample.

By selecting stars with accurate parallaxes from the Hipparcos catalog we introduced a distance bias which depends on the absolute magnitude: the stars which are intrinsically brighter can be accurately measured out to a larger distance. Therefore, there are relatively more bright stars in our Hipparcos sample than in our OGLE sample, as it is apparent in Fig.3. The difference in the slope of the red giant “background”, i.e. the different value of the ‘b’ coefficient in Eq.1, introduces a shift in the location of the gaussian peak describing the red clump. In order to correct for the differences in the two samples we multiplied the number of Hipparcos stars in each magnitude bin by the factor

$$f \sim \frac{a_{BW} + b_{BW}M_I}{a_H + b_H M_I}. \quad (2)$$

Following this procedure the new normalized fit to the modified Hipparcos histogram gave new values of the (a_H, b_H) coefficients consistent with their Baade’s Window values, and




Fig. 4.— The number of red clump stars in the solar neighborhood, based on Hipparcos data, is shown as a function of their $(V - I)$ color with the thin solid line. The number of red clump stars in Baade’s Window, based on the OGLE data, is shown as a function of their $(V - I)_0$ color with the thick solid line. All distributions are normalized, and include all stars with their absolute magnitude in the range: $-0.485 < M_I < 0.115$, i.e. within $0.3; mag$ of $M_{I,m}$. The bulge stars are much redder and presumably more metal rich than nearby disk stars.

also a new value of the gaussian peak: $M_{I,m} = -0.125 \pm 0.019$, which differs by 0.06 mag from our original estimate.

In order to estimate the effect of interstellar extinction we selected a sub-sample of 228 Hipparcos stars with the distance $d < 70$ pc, and we determined the parameters of the best fit to the luminosity function described with Eq.1, correcting for the (now reduced) distance bias in the way described in the previous paragraph. We found that $M_{I,m} = -0.192 \pm 0.023$, and $\sigma_{RC} = 0.208$ mag for these nearby stars, with the average distance $\langle d_{<70} \rangle = 50$ pc. For the remaining 437 Hipparcos red clump giants with the distance $d > 70$ pc we obtained $M_{I,m} = -0.094 \pm 0.027$, $\sigma_{RC} = 0.234$ mag and the average distance $\langle d_{>70} \rangle = 106$ pc. The more distant stars are fainter by 0.098 mag, with the difference in average distance being 56 pc. The most obvious explanation for this difference is the interstellar extinction. Extrapolating these results to the distance $d = 0$, and therefore to zero extinction, we obtain $M_{I_0,m} = -0.279$. Our final estimate of the correct average magnitude of the nearby

red clump stars is 0.088 mag brighter than our original crude estimate. We adopt this difference as our crude estimate of the possible systematic error in the absolute magnitude determination of the nearby red clump giants.

The centroid of the gaussian distribution of red clump giants in Baade’s Window corresponds to the distance which differs from the galactocentric distance for two reasons. First, because of bar asymmetry the point with the highest space density of the red clump stars seen in the direction $(l, b) = (1, -4)$ deg is somewhat closer than the galactic center. Second, the volume within the observed solid angle increases with the distance, which implies that we see more stars on the far side than on the near side of the bar, and the observed centroid is shifted to a somewhat larger distance than the location of the highest space density of the bar stars. We used the E2 bar model of Stanek et al. (1997) to calculate the two opposing effects, and we found that the galactocentric distance modulus is larger by 0.02 mag than the distance modulus obtained by us for the apparent centroid of the star distribution in Baade’s Window. Therefore, the apparent magnitude of red clump giants at the galactic center distance, corrected for reddening, is $I_{0,m} = 14.343$ mag, with the error due to our model estimated at 0.02 mag. A larger contribution to the error comes from zero point in the interstellar extinction, estimated to be ± 0.05 mag (Gould, Popowski & Terndrup 1997; Alcock et al. 1997).

Combining the corrected values of the absolute magnitudes of the nearby red clump stars, $M_{I_{0,m}} = -0.279 \pm 0.088$, with the apparent magnitude of the red clump stars at the galactocentric distance, $I_{0,m} = 14.343 \pm 0.05$, we obtain the corrected distance modulus to the galactic center: $(M - m)_{GC} = 14.62 \pm 0.10$, and the corresponding distance $R_0 = 8.4 \pm 0.4$ kpc.

There is yet another type of a systematic error possible: the age, the chemical composition, and the masses of red clump giants may be systematically different in the bulge/bar and near the sun. The apparent systematic difference in the colors implies that the two populations are in fact different. Recent stellar evolutionary models (Jimenez et al. 1997) indicate that in the age range 2 – 12 billion years the effective temperature is dominated by the metallicity. This suggests that the bulge population is on average more metal rich than the solar region. This seems to be in conflict with McWilliam & Rich (1994), who found that giants in the bulge and locally have similar metallicity. It also seems that among the bulge and local giants the most extreme stars are about equally super metal-rich (Castro et al. 1996).

It is not clear at this time what is the reason for the major difference in colors between the bulge and the local clump giants, but at least two possibilities may be considered. The spectroscopic samples are relatively small, and the selection criteria are not clear. On the

other hand, the photometric samples are more complete, or at least they should have no color bias. It is possible, that while the range of metallicity in the bulge and near the sun is similar, the distributions are different, with the super metal-rich stars more common in the bulge. Another possibility is that photometry and spectroscopy refer to different metallicities. The $(V - I)$ color is determined mostly by the effective temperature, which in turn depends on the photospheric continuum opacity in stars with deep convective envelopes, such as the red clump giants. It is not obvious that the spectroscopic metallicity based on line opacities is weighted in the same way as the photometric metallicity based on continuum opacity. In any case, the issue of color difference has to be resolved for the distance estimate based on the clump giants to be trusted.

4. Discussion

We found empirically that the average absolute magnitude of the red clump stars as measured in the Cousins’s I-band is independent on their $(V - I)$ color. It is important to reproduce this property with theoretical models, to have a better assessment of the robustness of our assertion, i.e. to have theoretical values of M_I calculated for a broad range of stellar ages and chemical compositions. Note that stellar populations are different in the galactic bulge and in the solar neighborhood, and a theoretical understanding of the $M_{I,m} - (V - I)$ relation is highly desirable.

Among the various stellar distance indicators the red clump giants are likely to be the best for determining the galactocentric distance because there are so many of them. In particular Hipparcos provided accurate distance determinations for almost 2,000 such stars, but unfortunately I-band photometry is available for only $\sim 30\%$ of them. It is important to obtain I-band photometry for all Hipparcos red clump giants in order to make a better correction for at least two effects: the distance bias in the catalog and the interstellar extinction. It would be useful to make accurate I-band photometry for all $\sim 10^6$ stars in the Tycho catalogue (Høg et al. 1997). Another important work which will be done soon with the forthcoming OGLE-2 BVI photometry of $\sim 3 \times 10^7$ stars in the bulge (Udalski, Kubiak & Szymański 1997), is a better bulge/bar model and a better extinction map of that region. This will reduce the uncertainty in the OGLE/Hipparcos comparison.

It is a great pleasure to acknowledge helpful suggestions by Andy Gould, Michael Richmond, Michael Strauss and Przemek Wozniak, and the anonymous referee. This work was supported by the NSF grants AST–9313620 and AST–9530478. KZS was supported by the Harvard-Smithsonian Center for Astrophysics Fellowship.

REFERENCES

- Alcock, Ch. et al. 1997, ApJ, submitted (astro-ph/9706292)
- Castellani, V., Chieffi, A., & Straniero, O. 1992, ApJS, 78, 517
- Castro, S., Rich, R. M., McWilliam, A., Ho, L. C., Spinrad, H., Filippenko, A. V., & Bell, R. A. 1996, AJ, 111, 2439
- Gould, A., Popowski, P., & Terndrup, D. M. 1997, ApJ, submitted (astro-ph/9705020)
- Høg, E. et al. 1997, A&A, 323, L57
- Jimenez, R., Flynn, C., & Kotoneva, E. 1997, MNRAS, submitted (astro-ph/9709056)
- Kiraga, M., Paczyński, B., & Stanek, K. Z. 1997, ApJ, 485, 611
- McWilliam, A., & Rich, R. M. 1994, ApJS, 91, 749
- Paczynski, B. et al. 1994, AJ, 107, 2060
- Perryman, M. A. C. et al. 1997, A&A, 323, L49
- Reid, M. J. 1993, ARA&A, 31, 345
- Seidel, E., Demarque, P., & Weinberg, D. 1987, ApJS, 63, 917
- Stanek, K. Z., et al. 1994, ApJ, 429, L73
- Stanek, K. Z. 1995, ApJ, 441, L29
- Stanek, K. Z. 1996, ApJ, 460, L37
- Stanek, K. Z., et al. 1997, ApJ, 477, 163
- Udalski, A., et al. 1992, AcA, 42, 253
- Udalski, A., et al. 1993, AcA, 43, 69
- Udalski, A., Kubiak, M., & Szymański, M., 1997, AcA, 47, 319 (astro-ph/9710091)
- Woźniak, P. R., & Stanek, K. Z. 1996, ApJ, 464, 233